

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Density Distribution of the Warm Ionized Medium

Alex S. Hill¹, Ronald J. Reynolds¹, Robert A. Benjamin², and L. Matthew Haffner¹

¹*University of Wisconsin-Madison (hill@astro.wisc.edu),* ²*University of Wisconsin-Whitewater*

Abstract. Observations of H α emission measures and pulsar dispersion measures at high Galactic latitude ($|b| > 10^\circ$) provide information about the density and distribution of the diffuse warm ionized medium (WIM). The diffuse WIM has a lognormal distribution of $\text{EM} \sin |b|$, which is consistent with a density structure established by isothermal turbulence. The H⁺ responsible for most of the emission along high-EM $\sin |b|$ sightlines is clumped in high density ($> 0.1 \text{ cm}^{-3}$) regions that occupy only a few parsecs along the line of sight, while the H⁺ along low-EM sightlines occupies hundreds of parsecs with considerably lower densities.

1. Introduction

The warm ionized medium (WIM) is a major component of our Galaxy. The WIM consists of a pervasive, diffuse plasma layer with a temperature $T \sim 8000 \text{ K}$, a scale height of $h \approx 1 \text{ kpc}$ (more than 3 times the scale height of neutral hydrogen), and densities of $\approx 0.08 \text{ cm}^{-3}$ which occupy $\approx 20\%$ of the volume of the disk (Reynolds 1991). The ionization and heating sources of the WIM and its relationship to the other phases of the ISM are not well understood.

We use observations of interstellar H α emission and pulsar dispersion measures to explore the density and distribution of diffuse plasma in the WIM. Pulsar dispersion measures provide the column density, $\text{DM} = \int_0^D n_e dl$, of the free electrons along the line of sight to the pulsar. H α emission results from recombination in ionized regions; it is proportional to the square of the electron density along the entire line of sight: $\text{EM} = \int_0^\infty n_e^2 dl$.

2. H α data

The Wisconsin H-Alpha Mapper (WHAM) Northern Sky Survey (WHAM-NSS; Haffner et al. 2003) mapped the H α ($\lambda 6563$) optical emission line in the northern sky with a spectral resolution of 12 km s^{-1} and an angular resolution of 1° . We use a restricted set of the WHAM-NSS northern sky map integrated over all velocities. We have removed lines of sight that intersect classical H II regions, leaving a sample of only the diffuse WIM (Figure 1).




Figure 1. Map of $\text{EM} \sin |b|$ from the WHAM Northern Sky Survey for Galactic latitude $|b| > 10^\circ$ with sightlines that intersect known classical H II regions eliminated. Dashed lines represent 30° increments in longitude and 15° increments in latitude.

3. Distribution of $\text{H}\alpha$ emission

A histogram of $\text{EM} \sin |b|$, the component of emission measure perpendicular to the Galactic plane, for $|b| > 10^\circ$ is shown in Figure 2. We fit a normal distribution to the data (left panel of Figure 2). The sharp peak indicates that the emission measure along most sightlines is dominated by plasma in a disk-like distribution having a mean value of $\text{EM} \sin |b|$ of $1.1 \pm 0.5 \text{ pc cm}^{-6}$. There are considerable enhancements above a normal distribution at high $\text{EM} \sin |b|$ ($\geq 2 \text{ pc cm}^{-6}$), indicating the presence of high- $\text{EM} \sin |b|$ regions along about 1/5 of sightlines.

The entire diffuse WIM, including the enhanced high-EM tail, is fit well by a lognormal distribution in $\text{EM} \sin |b|$ or, equivalently, a Gaussian distribution in $\log \text{EM} \sin |b|$ (right panel of Figure 2). The classical H II regions (not shown) are also well fit by a lognormal distribution with different parameters. A lognormal distribution is the expected probability density function in density for a plasma with a density structure established by isothermal turbulence (e. g., Vázquez-Semadeni & Passot 1999). We are currently examining how a lognormal distribution in EM can be used to diagnose the turbulence in the WIM.




Figure 2. Histograms of $\text{EM sin } |b|$ (*left*) and $\log \text{EM sin } |b|$ (*right*) for diffuse WIM sightlines in the WHAM-NSS survey (see text). A Gaussian fits the peak of the distribution in linear space well (dashed line), but the enhanced tail at $\text{EM sin } |b| > 2 \text{ pc cm}^{-6}$ is typical of a lognormal distribution (dotted line, *right*).

4. Density and occupation length of the diffuse gas

Comparisons of pulsar dispersion measures and $\text{H}\alpha$ emission measures along the same sightline provide information about the distribution of the gas which contributes to the EM and DM. There are 194 pulsars along diffuse WIM sightlines in the WHAM-NSS region (Manchester et al. 2005). For high-EM sightlines ($\text{EM sin } |b| > 2 \text{ pc cm}^{-6}$), the mean dispersion measure is $\langle \text{DM sin } |b| \rangle = 14.8 \pm 0.9 \text{ pc cm}^{-3}$, whereas for low-EM sightlines ($\text{EM sin } |b| < 2 \text{ pc cm}^{-6}$), the mean value is $\langle \text{DM sin } |b| \rangle = 13.5 \pm 0.5 \text{ pc cm}^{-3}$. Thus, the fluctuations in density that dominate the emission measure along the high-EM sightlines contribute little to the total H^+ column density.

The characteristic electron density n_e and occupation length L_e of an ionized region can be estimated from $n_e = \text{EM}/\text{DM}$ and $L_e = (\text{DM})^2/\text{EM}$ (Reynolds 1991). Figure 3 shows the relationship between the density and size of ionized regions. The gas along sightlines with densities $n_e \sim 0.1 \text{ cm}^{-3}$ typically occupies a few hundred parsecs along the line of sight, whereas the high-EM gas has densities approaching 1 cm^{-3} that occupy only a few parsecs along the line of sight.

The local electron density and occupation length of the diffuse H^+ can be determined toward pulsars in globular clusters more than 3 kpc above the Galactic midplane, where DM samples the total H^+ column density through the disk. Eight such globular clusters are known to contain pulsars, but two are outside the WHAM-NSS survey region. Data for these globular clusters are shown in Table 1.




Figure 3. Occupation length L_c versus characteristic density n_c for gas along 194 pulsar sightlines which are in the WHAM-NSS survey area and not towards classical H II regions.

5. Summary

We are investigating the density and distribution of the diffuse plasma in the warm ionized medium. The H α emission from the WIM follows a lognormal distribution in $EM \sin |b|$. A lognormal distribution in density is typical of isothermal turbulence, but its significance for integrated density (EM and DM) is not yet clear. Pulsar dispersion measure data indicate that localized (a few parsecs in length), high-density ($> 0.1 \text{ cm}^{-3}$) regions dominate the emission measure at the high end of the EM distribution while contributing little to the total DM.

References

- Haffner, L. M., Reynolds, R. J., Tufte, S. L., Madsen, G. J., Jaehnig, K. P., & Percival, J. W. 2003, *ApJS*, 149, 405

Globular Cluster	DM sin $ b $ (pc cm ⁻³)	EM sin $ b $ (pc cm ⁻⁶)	n_c (cm ⁻³)	L_c sin $ b $ (pc)
M3	25.888	0.88	0.034	760
M5	21.796	1.3	0.061	360
M13	19.577	1.0	0.051	380
M15	30.684	3.2	0.10	300
M30	18.283	1.7	0.094	190
M53	23.621	1.1	0.046	520

Table 1. Properties of lines of sight towards $|z| > 3$ kpc globular clusters with known pulsars. EM data from WHAM-NSS; DM data from the ATNF pulsar catalog (Manchester et al. 2005). Sightlines towards M3 and M5 are contaminated by bright stars, so we used an average of nearby sightlines to determine the EM.

Manchester, R. N., Hobbs, G. B., Teoh, A., & Hobbs, M. 2005, AJ, 129, 1993

Reynolds, R. J. 1991, ApJ, 372, L17

Vázquez-Semadeni, E. & Passot, T. 1999, in *Interstellar Turbulence*, ed. J. Franco & A. Carramiñana (Cambridge University Press), 223